

For Your Information

Courtesy Of:

Cotati Large Animal Hospital

8079 Gravenstein Hwy, Cotati, CA 94931

(707) 795-4356 (707) 578-4760

Gene Harlan, DVM Calvin Dotti, DVM Andrew Wirz, DVM Sally Venable, DVM

www.cotatilargeanimal.com

Dairy Cattle Vaccine Recommendations

- **Newborn calf or when purchased**
 - 2cc Nasalgen in the nostril
- **3-4 months of age**
 - Dehorn
 - 2cc Vision 8 (Sub Q)
 - 2cc Nasalgen in the nostril
 - 5cc Triangle 10 (Sub Q)
 - 2cc Pinkeye (Sub Q)
 - Brucellosis (**must** be given by the veterinarian between 4-12 months old)
 - Deworm (not organic)
 - Pour on fly protection (not organic)
- **Booster vaccinations (4-8 weeks after the 1st set of vaccines)**
 - 2cc Vision 8 (Sub Q)
 - 2cc Nasalgen in the nostril
 - 5cc Triangle 10 (Sub Q)
 - 2cc Pinkeye (Sub Q)
 - Deworm (not organic)
 - Pour on fly protection (not organic)
- **10-12 months of age**
 - 2cc Vision 8 (Sub Q)
 - 2cc Nasalgen in the nostril
 - Bovishield FB 5L5 HB (Sub Q)
 - Or 2cc Triangle 5 (Sub Q)
 - And Spirovac VL5 (Sub Q)
 - If breeding with bull instead of AI
 - Deworm (not organic)
 - Magnet
- **Cows annually**
 - 5cc Triangle 10 (Sub Q)
 - 2cc Vision 8 (Sub Q)
 - 5cc Trivib 5L (Sub Q) at pregnancy check