

For Your Information

Courtesy Of:

Cotati Large Animal Hospital

8079 Gravenstein Hwy, Cotati, CA 94931

(707) 795-4356 (707) 578-4760

Gene Harlan, DVM Calvin Dotti, DVM Andrew Wirz, DVM Sally Venable, DVM

www.cotatilargeanimal.com

BEEF CATTLE

- I. **Newborn**
 - a. Inforce - first week
- II. **2-3 Months of Age**
 - a. Nasalgen or Inforce (2cc IN)
 - b. Triangle-10 (5cc SQ)
 - c. Vision-8 (2cc SQ) or Covexin 8 (5cc SQ) (Clostridium)
 - d. Ivomec wormer (1cc / 100 lbs) or Cydectin (1cc / 110lbs)
 - e. Dehorn & castrate bulls
 - i. Tetanus Antitoxin
 - f. Fly & lice control (Permethrin CDS or Cylence)
- III. **4-6 Months of Age (1 Month later)**
 - a. Nasalgen
 - b. Triangle-10
 - c. Vision-8 or Covexin 8 (Clostridium)
 - d. Panacur (2.5cc/ 100 lbs) or Valbazen wormer (1cc / 25 lbs)
 - e. Fly & lice control
 - f. Brucellosis – **heifers only**
- IV. **Replacement Heifers – Weaning**
 - a. Bovishield 5PFL5HB (2cc SQ) or Triangle-10
 - b. Clostridium
 - c. Nasalgen
 - d. Wormer
 - e. Fly & lice control
- V. **Cow Herd & Heifers Before Breeding – Fall**
 - a. Triangle-10
 - b. Clostridium
 - c. Ivomec wormer or Cydectin
 - d. Fly & lice control
- VI. **Cow Herd – Spring**
 - a. Triangle-5 (2cc SQ)
 - b. Spirovac VL5 (5cc SQ) (Booster 1st year) or TriVib5L (5cc SQ)
 - c. Clostridium
 - d. Panacur wormer
 - e. Fly & lice control
- VII. **Bulls**
 - a. Breeding soundness exam 1-2 months before breeding season and Trichomonosis test
 - b. Triangle-5
 - c. Clostridium
 - d. Trichguard (2 injections if possible before breeding) (2cc SQ)
 - e. TriVib5L
 - f. Ivomec/Cydectin/Valbazen/or Panacur Wormer
 - g. Fly & Lice control
 - h. Anaplasmosis (optional) (1cc SQ)